

AcrySof® IQ PanOptix® IOL

The next-generation trifocal from the global leaders in eye care

What are cataracts and how are they treated?

Cataracts occur when the lens of the eye becomes cloudy with age.¹ Symptoms include:

- Blurry vision
- Dull, yellowish colours
- Glare from lights
- Poor night vision¹

Cataracts are treated with surgery to replace the old lens with an artificial one called an Intraocular lens (IOL).¹

What is AcrySof® IQ PanOptix® IOL?

AcrySof° IQ PanOptix° IOL is an intraocular lens designed to provide you with clear vision for near (40 cm), intermediate (60 cm), and far distances without glasses. Because AcrySof° IQ PanOptix° IOL helps with all 3 distances, it is sometimes called a "trifocal". ²

Why has my eye surgeon recommended AcrySof® IQ PanOptix® IOL?

Your surgeon has recommended the AcrySof° IQ PanOptix° IOL lens to reduce or eliminate your need for glasses after cataract surgery at near, intermediate and far distances.^{2,3}

After receiving AcrySof® IQ PanOptix® IOL, will I have the vision I had in my thirties before reading glasses?


There is no intraocular lens that exactly mimics a young eye. But trifocal lenses offer the latest advances in vision technology. They are designed to enhance your lifestyle by providing clear vision at near, intermediate and far distances.^{2,3}

Will having AcrySof® IQ PanOptix® IOL mean I never have to wear glasses or contact lenses again?

AcrySof° IQ PanOptix° IOL is designed to provide good vision in most day-to-day situations without the need for glasses.² You may find that your vision for a few detailed tasks (e.g., reading small print) can be optimised further by wearing glasses. This is common to all intraocular lenses.²







After surgery What can I expect from my AcrySof® IQ PanOptix® IOL?

Day 1-2 after surgery

Ask your eye surgeon how to care for your eye after surgery, when you can return to your usual activities, and what to watch for (including signs that you should contact your surgeon).

You will have a follow-up appointment a day or two after your surgery⁷

- If surgery is on your first eye, your vision might not feel completely clear until your second eye is treated.
- If this is your second eye, most people find they have good vision at near, intermediate and far distances within a few days of surgery.

Things you may notice in the first day or two:

- Some glare with bright lights
- Halos around lights, especially at night²

Many people barely notice the glare and halos at all or feel they can ignore them. Some people notice them all the time, and a few are significantly bothered by

Your brain is generally able to adapt and the glare and halos will become less noticeable over time.7

This picture demonstrates what glare (streetlights and car headlights) and halos (road sign and white stripe on the edge of the road) might look like.



(40 cm)

Year 1 after surgery and beyond It is unlikely your vision will change significantly in the years after your surgery. Have your vision and eye health tested: • at least every 2 years if you are under 65 at least annually if you are 65 and over • if you notice any changes in your vision 8 Intermediate vision (60 cm) **Far vision**

Doctor Discussion Guide

Use this guide to help you prepare for your next appointment

Visit **PanOptix.ca** to learn more about AcrySof® IQ PanOptix® IOL.

Getting ready for your appointment:

- 1. Whenever you think of a question to ask, write it down on this guide.
- 2. Consider bringing a friend or family member to your appointment.
- 3. Bring a pen or pencil and use it to take notes on this guide.
- 4. Leave yourself plenty of travel time to reach your appointment.

Questions for your eye surgeon:

- 1. Why have you recommended the AcrySof° IQ PanOptix° IOL for me?
- 2. What will my vision be like after the surgery?
- 3. Will I still need to wear glasses or contacts after the surgery?
- 4. What are the risks of surgery?
- 5. How long does the surgery take?
- 6. How should I prepare for the surgery (e.g., do I need to stop any of my medications)?
- 7. How do I care for myself after the surgery?
- 8. When can I return to my usual activities?
- 9. What are the signs of complications, and what should I do if they occur?

10. Other:			
11. Other:			
12. Other:			

Notes:				
Surgeon nar	me and contac	ct informatio	n:	
Next appoin	tment:			

AcrySof® IQ PanOptix® Presbyopia Correcting Intraocular Lenses Important Product Information

INDICATIONS:

The AcrySof° IQ PanOptix° Presbyopia Correcting Intraocular Lens is intended for primary implantation in the capsular bag in the posterior chamber for the visual correction of aphakia secondary to removal of a cataractous lens in adult patients with and without presbyopia, who desire near, intermediate and distance vision with increased spectacle independence.

WARNINGS:

As with any surgical procedure, there are associated risks. Careful preoperative evaluation and sound clinical judgment should be used by the surgeon to decide the risk/benefit ratio before implanting a lens of this type. This is particularly so in a patient with any of the conditions described in the AcrySof 'IQ PanOptix' physician labeling. Some patients may experience visual disturbances and/or difficulty seeing due to the multifocal lens design, especially under dim light conditions. As with other multifocal IOLs, visual symptoms may be significant enough that the patient will request explantation of an AcrySof' IQ PanOptix' IOL. Posterior capsule opacification (PCO) may significantly affect the vision of patients with multifocal IOLs sooner in its progression than in patients with monofocal IOLs.

ATTENTION:

Refer to the Directions for Use labeling for Model TFNT00 for a complete listing of indications, warnings and precautions.

REFERENCES:

- 1. Canadian Ophthalmological Society. Cataract. http://www.cos-sco.ca/vision-health-information/conditions-disorders-trea*ents/cataract/.
- 2. AcrySof° IQ PanOptix° IOL Directions for Use.
- 3. Alcon Data on File. TDOC-0053542 (Mar 10 2017).
- 4. Canadian National Institute for the Blind. Light up your life. http://www.cnib.ca/en/your-eyes/eye-care/lighting-0607/Pages/default.aspx.
- 5. Charness N, Dijkstra K, Jastrzembski T, et al. Monitor viewing distract for younger and older workers. Proceedings of the Human Factors and Ergonomics Society 52nd Annual Meeting, 2008. http://www.academia.edu/477435/Monitor_Viewing_Distance_for_Younger_and_Older_Workers. Accessed May 9, 2015.
- 6. Kasetsuwan N, Satitpitakul V, Changul , Jariyakosol S. Incidence and pattern of dry eye after cataract surgery. PLos ONE 2013;8(11):e78657. http://journals.plos. org/plosone/article/file?id=10.1371/journal.pone.0078657&type=printable.
- 7. Canadian Ophthalmological Society. Canadian Ophthalmological Society evidence-based clinical practice guidelines for cataract surgery in the adult eye. Can J Ophthalmol. 2008;43(Suppl 1): S1-S57.
- 8. Canadian Association of Optometrists. The eye exam. https://opto.ca/health-library/the-eye-exam. Accessed May 18, 2017.



